

AD-A072 226

MIDWEST RESEARCH INST KANSAS CITY MO
CONVERGENCE PROOF OF ECONOMIC REPRESENTATION OF
JUL 60 J L FIELDS, Y L LUKE

F/G 12/1
--ETC(U)
NONR-2638(00)

UNCLASSIFIED

NL


| OF |
AD
A072.26

DATE
FILED



END
DATE
FILMED
9-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL

COLUMBIA UNIVERSITY
HUDSON LABORATORIES
CONTRACT NO-DNR-27135

CONVERGENCE PROOF OF ECONOMIC REPRESENTATION
OF TRANSCENDENTAL FUNCTIONS

1

TECHNICAL REPORT
July 15, 1960


M.R.I. Project No. 2229-P

Contract No. Nonr-2638(00)(X)

DDC FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited



For

Applied Mathematics Laboratory
David W. Taylor Model Basin
Washington, D. C.

78 03 07 094

79 08 03 117 AUG 10 1960

MIDWEST RESEARCH INSTITUTE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CONVERGENCE PROOF OF ECONOMIC REPRESENTATION OF TRANSCENDENTAL FUNCTIONS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Fields, Jerry L. and Luke, Yudell L.		8. CONTRACT OR GRANT NUMBER(s) N0nr-2638(00)(X)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Midwest Research Institute		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 2229-P
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 220 800 North Quincy Street Arlington, VA 22217		12. REPORT DATE 15 JUL 1960
		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

✓ MIDWEST RESEARCH INSTITUTE

COLUMBIA UNIVERSITY
HUDSON LABORATORIES
CONTRACT N6-ONR-27135

⑥
CONVERGENCE PROOF OF ECONOMIC REPRESENTATION
OF TRANSCENDENTAL FUNCTIONS

by

⑩ Jerry L. Fields
Yudell L. Luke

⑪ 15 Jul 60

⑫ 31 p.

⑨
TECHNICAL REPORT

July 18, 1960

M.R.I. Project No. 2229-P

✓ Contract No. ⑬ Nonr-2638(00)(X)

For

Applied Mathematics Laboratory
David W. Taylor Model Basin
Washington, D. C.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

79 08 03 117
230350
78 03 07 094

AUG 10 1960

PREFACE


This report covers research initiated by the Applied Mathematics Laboratory, David W. Taylor Model Basin, Washington, D. C. The research work upon which this report is based was accomplished by Midwest Research Institute under Contract No. Nonr-2638(00)(X).

The numerics in this report are related to work previously done in earlier reports, and the authors wish to take this opportunity to again thank those members of the Midwest Research Institute staff who materially contributed. They are Geraldine Coombs, Betty Kahn, Anna Lee Samuels and Wanda Shelp.

J.L.F. and Y.L.L.

Approved for:

MIDWEST RESEARCH INSTITUTE



S. L. Levy, Manager
Mathematics and Physics Division

July 15, 1960

TABLE OF CONTENTS

	<u>Page No.</u>
Summary and Introduction	1
1. Convergence of the Rational Approximations	2
2. Numerics	13
Bibliography	16

CONVERGENCE PROOF OF ECONOMIC REPRESENTATION
OF TRANSCENDENTAL FUNCTIONS

by

Jerry L. Fields

and

Yudell L. Luke

Summary and Introduction

In a previous report [1]^{1/}, it was shown that a large class of transcendental functions could be represented by a rational function, the ratio of two polynomials, together with a remainder term. More specifically, if $E(z)$ is defined by a Laplace integral, then

$$E(z) = \phi_n/f_n + R_n, \quad R_n = F_n/f_n$$

where ϕ_n and F_n are polynomials of degree n and R_n is the remainder. In Section I it is shown that this can be considered a regular summation technique.

By proper choice of some parameters, f_n is a polynomial of hypergeometric form. If, except for a multiplicative factor, $E(z)$ has a generalized hypergeometric series representation, then F_n is a sum of n generalized hypergeometric series.

Only in certain cases could it be shown that the rational representation converges, i.e., for z fixed, $\lim_{n \rightarrow \infty} R_n(z) = 0$. One of the principal difficulties lies in the fact that asymptotic expansions of a certain class of hypergeometric polynomials were known only for a few special cases, the classical Jacobi polynomials, for example. Asymptotic expansions of this class of hypergeometric polynomials now exist, see [2] and the convergence question can be further explored.

^{1/} Numbers in square brackets pertain to references at end of report.

The structure of the numerator of the error term, i.e., F_n is complicated and only in a few special cases has it been possible to represent it in a useful form. This drawback is partly alleviated by the fact that one of the above-mentioned special cases includes the Whittaker functions and their natural generalizations. For those cases where F_n can be put into a useful form, we prove convergence of the rational representation.

In the case of particular interest, i.e., the Whittaker functions, we represent F_n as an integral and then find a suitable bound for it. This coupled with an asymptotic estimate of f_n enables us to prove that $\lim_{n \rightarrow \infty} R_n = 0$ and also to obtain an asymptotic bound for the error. This is the essence of Section I. In Section II, numerics are presented to display the effectiveness of this bound.

1. Convergence of the Rational Approximations

In [1], it was shown how to formally generate a rather general sequence of rational approximations for the generalized hypergeometric function [3]

$$E(z) = {}_{p+1}F_q \left(\begin{matrix} \sigma, \alpha_1, \dots, \alpha_p \\ \beta_1, \dots, \beta_q \end{matrix} \middle| \frac{\lambda}{z} \right), \quad (1.1)$$

where

$${}_mF_n(x) = {}_mF_n \left(\begin{matrix} \alpha_1, \dots, \alpha_m \\ \beta_1, \dots, \beta_n \end{matrix} \middle| x \right) = \sum_{k=0}^{\infty} \frac{\prod_{t=1}^m (\alpha_t)_k}{\prod_{t=1}^n (\beta_t)_k} \frac{(x)^k}{k!} \quad (1.2)$$

$$(\omega)_k = \frac{\Gamma(\omega+k)}{\Gamma(\omega)}$$

Throughout this paper we employ a contracted notation and write

$${}_m F_n(x) = {}_m F_n \left(\begin{matrix} \alpha_m \\ \rho_n \end{matrix} \middle| x \right) = \sum_{k=0}^{\infty} \frac{(\alpha_m)_k}{(\rho_n)_k} \frac{(x)^k}{k!} \quad (1.3)$$

Thus $(\alpha_m)_k$ is interpreted as $\prod_{t=1}^m (\alpha_t)_k$, and a similar remark holds for $(\rho_n)_k$. We assume that none of the α_t 's, or ρ_t 's are zero or a negative integer. Also, it is assumed that the difference of any numerator parameter α_j , and any denominator parameter ρ_t , is never equal to zero. Empty terms in expression such as (1.3) are replaced by unity. Two distinct sequences of rational approximations were developed in the above paper [1]. They are as follows.

Type I or Homogeneous Case

$$E(z) = \frac{\phi_n(z, \gamma)}{f_n(\gamma)} + R_n(z, \gamma), \quad R_n(z, \gamma) = \frac{F_n(a_k, z, \gamma)}{f(\gamma)} \quad (1.4)$$

$$\phi_n(z, \gamma) = \sum_{r=0}^n \frac{(\sigma^-)_r (\alpha_p)_r}{(1)_r (\rho_q)_r} \left(\frac{\lambda}{z} \right)^r f_n^{[r]}(\gamma) \quad (1.5)$$

$$f_n^{[r]}(\gamma) = \sum_{k=r}^n a_{n-k} \gamma^k, \quad f_n^{[0]} = f_n(\gamma) \quad (1.6)$$

$$F_n(a_k, z, \gamma) = \frac{z^\sigma}{\Gamma(\sigma)} \int_0^\infty e^{-zt} t^{\sigma-1} \sum_{k=0}^n a_{n-k} (\lambda \gamma t)^k \sum_{r=0}^\infty \frac{(\alpha_p)_{r+k+1} (\lambda t)^{r+1}}{(\rho_q)_{r+k+1} (r+k+1)!} dt \quad (1.7)$$

$$= \frac{z^\sigma}{\Gamma(\sigma)} \int_0^\infty e^{-zt} t^{\sigma-1} \sum_{k=0}^n a_{n-k} \left(\frac{\lambda \gamma}{z} \right)^k \sum_{r=0}^\infty \frac{(\alpha_p)_{r+k+1} (\sigma+r+1)_k (\lambda t)^{r+1}}{(\rho_q)_{r+k+1} (r+k+1)!} dt \quad (1.8)$$

$$= \frac{\sigma \lambda \alpha_p}{z \rho_q} \sum_{r=0}^\infty \frac{(\sigma+1)_r (\alpha_p+1)_r}{(2)_r (\rho_q+1)_r} \left(\frac{\lambda}{z} \right)^r \sum_{k=0}^n \frac{a_{n-k} (\sigma+r+1)_k (\alpha_p+r+1)_k}{(2+r)_k (\rho_q+r+1)_k} \left(\frac{\lambda \gamma}{z} \right)^k \quad (1.9)$$

where $\alpha_p = \prod_{t=1}^p \alpha_t$, and $\rho_q = \prod_{t=1}^q \rho_t$. Here the coefficients a_k are as yet arbitrary, and γ is unrestricted except that $|\gamma/z| \leq 1$.

Type II or Non-Homogeneous

$$E(z) = \frac{\Phi_n(z, \gamma)}{f_n(\gamma)} + R_n(z, \gamma), \quad R_n(z, \gamma) = \frac{F_n(a_k, z, \gamma)}{f_n(\gamma)} \quad (1.10)$$

$$\Phi_n(z, \gamma) = \sum_{r=0}^{n-1} \frac{(\sigma)_r (\alpha_p)_r}{(1)_r (\rho_q)_r} \left(\frac{\lambda}{z} \right)^r f_n^{[r+1]}(\gamma) \quad (1.11)$$

$$f_n^{[r]}(\gamma) = \sum_{k=r}^n a_{n-k} \gamma^k, \quad f_n^{[0]}(\gamma) = f(\gamma) \quad (1.12)$$

$$F_n(a, z, \gamma) = \frac{z^\sigma}{\Gamma(\sigma)} \int_0^\infty e^{-zt} t^{\sigma-1} \sum_{k=0}^n a_{n-k} (\lambda \gamma t)^k \sum_{r=0}^\infty \frac{(\alpha_p)_{r+k} (\lambda t)^r}{(\rho_q)_{r+k} (r+k)!} dt \quad (1.13)$$

$$= \frac{z^\sigma}{\Gamma(\sigma)} \int_0^\infty e^{-zt} t^{\sigma-1} \sum_{k=0}^n a_{n-k} \left(\frac{\lambda \gamma}{z} \right)^k \sum_{r=0}^\infty \frac{(\alpha_p)_{r+k} (\sigma+r)_k (\lambda t)^r}{(\rho_q)_{r+k} (r+k)!} dt \quad (1.14)$$

$$= \sum_{r=0}^\infty \frac{(\sigma)_r (\alpha_p)_r}{(1)_r (\rho_q)_r} \left(\frac{\lambda}{z} \right)^r \sum_{k=0}^n \frac{a_{n-k} (\sigma+r)_k (\alpha_p+r)_k}{(1+r)_k (\rho_q+r)_k} \left(\frac{\lambda \gamma}{z} \right)^k \quad (1.15)$$

where, as before, the a_k 's are unspecified and γ is subject only to the restriction $|\gamma/z| \leq 1$.

The difference between the two sequences is that when $z = \gamma$, in the Type I representation, $\varphi_n(0) \neq 0$, whereas for the Type II representation under the same condition, $\varphi_n(0) = 0$. For some particular case one of the representations may be more desirable than the other. Also, it sometimes happens that one sequence approaches the desired limit monotonically from above, and the other monotonically from below, thus yielding rational inequalities for the functions in question. The above formal sequences were derived, strictly speaking, with the restrictions $p \leq q$ and $\text{Re}(z) > 0$ if $p < q$, while $\text{Re}(z) > \text{Re}(\lambda)$ if $p = q$. However, these restrictions may be weakened whenever the resulting expressions are defined and make sense.

The merit of [1] lies in the fact that it motivates how the arbitrary coefficients a_k should be chosen and gives a closed form expression for the error. The a_k 's were chosen in [1] to agree under certain circumstances, at least, with the rational approximations given by application of Lanczos' τ -method (see [4]) to the linear differential equation satisfied by $E(z)$ (see [3]). The choices made for the a_k were

Type I - Homogeneous

$$a_{n-k} = \frac{C_{nk} (\rho_q)_k}{(\alpha_p+1)_k \lambda^k}, \quad \text{if } \sigma = 1 \quad (1.16)$$

$$a_{n-k} = \frac{C_{nk}(\rho_q)_k(1)_k}{(\alpha_{p+1})_k(\sigma+1)_k \lambda^k}, \quad \text{if } \sigma \neq 1 \quad (1.17)$$

where

$$C_{nk} = \frac{(-n)_k(n+\alpha+\beta+1)_k}{(\beta+1)_k k!}.$$

Type II - Non-Homogeneous

$$a_{n-k} = \frac{C_{nk}(\rho_{q-1})_k(1)_k}{(\alpha_p)_k(\sigma)_k \lambda^k} \quad (1.18)$$

and C_{nk} is defined as above. With this selection of the coefficients a_k , we show that the rational sequences defined by (1.4) - (1.9) and (1.10) - (1.15) converge for two general classes of functions defined as follows: Class I is composed of those hypergeometric functions such that $p \leq q$ in (1.1) and Class II is made up of the hypergeometric functions such that $p = q+1$ in (1.1). Class II includes the Whittaker functions, modified Bessel functions, the Weber Parabolic Cylinder functions, and various special cases thereof. For completeness, we define a Class III composed of all hypergeometric functions such that $p \geq q+1$ in (1.1).

Considering $E(z)$ as a formal infinite series, we define the partial sum of this series by

$$P_n(z) = \sum_{r=0}^{n-1} \frac{(\sigma)_r(\alpha_p)_r}{(\rho_q)_r} \frac{(\lambda/z)^r}{r!}. \quad (1.19)$$

Rearrangement of (1.5) and (1.11) gives

$$\varphi_n(z, \gamma) = \sum_{k=a}^n a_{n-k} \gamma^k P_k(z) \quad (1.20)$$

where $a = 0$ for the Type I or Homogeneous Case,

$a = 1$ for the Type II or Non-Homogeneous Case.

Thus our rational approximations may be considered as a summation technique. A method of summability is said to be regular if it sums a convergent series to its ordinary sum. We now show that our economization process is a regular summability process.

Theorem. If

$$(1) \quad \varphi_n(z, \gamma) = \sum_{k=a}^n a_{n-k} \gamma^k P_k(z), \quad f_n(\gamma) = \sum_{k=0}^n a_{n-k} \gamma^k,$$

$$(2) \quad \lim_{k \rightarrow \infty} P_k(z) = E(z), \text{ for fixed } z,$$

$$(3) \quad a_{n-k} > 0, \quad \gamma > 0, \text{ and}$$

$$(4) \quad \lim_{n \rightarrow \infty} f_n(\gamma) = \infty, \quad f_{n+1}(\gamma) > f_n(\gamma) \text{ for fixed } \gamma,$$

$$\text{then } \lim_{n \rightarrow \infty} \left| \varphi_n(z, \gamma) / f_n(\gamma) - E(z) \right| = 0 \text{ for fixed } \gamma \text{ and } z.$$

Proof.

Given $\epsilon > 0$, there exists a positive integer N (usually dependent on z) such that $n > N$ implies $|P_k(z) - E(z)| < \epsilon/3$. Then

$$\begin{aligned} \varphi_n(z, \gamma) / f_n(\gamma) &= \sum_{k=a}^N \left\{ a_{n-k} \gamma^k [P_k(z) - E(z) - \epsilon/3] \right\} + \epsilon/3 \sum_{k=0}^N a_{n-k} \gamma^k \\ &\quad + \frac{\sum_{k=N}^n a_{n-k} \gamma^k [P_k(z) - E(z)] - a_n [\epsilon/3 + E(z)]}{f_n(\gamma)}. \end{aligned}$$

Thus, for $n > N$,

$$\left| \Phi_n(z, \gamma) / f_n(\gamma) - E(z) \right| \leq \frac{\sum_{k=a}^N \left\{ a_{n-k} \gamma^k |P_k(z) - E(z) - \epsilon/3| \right\} + a a_n (\epsilon/3 + |E(z)|)}{f_n(\gamma)} + \epsilon/3 \frac{f_n(\gamma)}{f_n(\gamma)}.$$

Since N is fixed, n can be chosen large enough such that the right hand side of the above is less than ϵ . This proves the theorem. From the results of [2], and (1.16) - (1.18), we see that convergence for Class I of the hypergeometric functions is proved, when $\lambda \geq 0$.

We now consider Class II, i.e., the situation where $p = q+1$. Treating the non-homogeneous or Type II case only, for $p = 1$ and $q = 0$, from (1.15) and (1.18) we get

$$F_n(z, \gamma) = \frac{z^\sigma}{\Gamma(\sigma)} \int_0^\infty e^{-zt} t^{\sigma-1} \sum_{k=0}^n C_{nk} (\gamma/z)^k {}_3F_2 \left(\begin{matrix} \alpha_1+k, \sigma+k, 1 \\ k+1, \sigma \end{matrix} \middle| \lambda t \right) dt. \quad (1.21)$$

By use of the relationships, see [3]

$${}_{p+1}F_{q+1} \left(\begin{matrix} \alpha_p, \sigma_1 \\ \rho_q, \sigma_1 + \sigma_2 \end{matrix} \middle| z \right) = \frac{\Gamma(\sigma_1 + \sigma_2)}{\Gamma(\sigma_1) \Gamma(\sigma_2)} \int_0^1 u^{\sigma_1-1} (1-u)^{\sigma_2-1} {}_pF_q \left(\begin{matrix} \alpha_p \\ \rho_q \end{matrix} \middle| zu \right) du, \quad \text{Re } \sigma_1 > 0, \text{ Re } \sigma_2 > 0 \quad (1.22)$$

and

$${}_2F_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| z \right) = (1-z)^{-\beta} {}_2F_1 \left(\begin{matrix} \gamma-\alpha, \beta \\ \gamma \end{matrix} \middle| \frac{z}{z-1} \right), \quad (1.23)$$

we can write

$$F_n(z, \gamma) = \frac{z^\sigma}{\Gamma(\sigma-1)\Gamma(1-\alpha_1)\Gamma(\alpha_1)} \int_0^\infty \int_0^1 \int_0^1 \frac{e^{-zt} t^{\sigma-1} u^{\alpha_1-1} (1-u)^{-\alpha_1} v^{\sigma-2}}{(1-t\lambda u)} dv du dt$$

$$\times {}_3F_2 \left(\begin{matrix} -n, n+\alpha+\beta+1, 1 \\ \alpha_1, 1+\beta \end{matrix} \middle| u \left(\frac{\gamma}{z} \right) \left(\frac{1-t\lambda uv}{1-t\lambda u} \right) \right) dv du dt ,$$

$$\operatorname{Re} \lambda < 0, \operatorname{Re}(z) > 0, 0 < \operatorname{Re} \alpha_1 < 1, \operatorname{Re} \sigma > 1 . \quad (1.24)$$

Defining

$$M_1^{II} = \max_{0 \leq w \leq 1} \left| {}_3F_2 \left(\begin{matrix} -n, n+\alpha+\beta+1, 1 \\ \beta+1, \alpha_1 \end{matrix} \middle| w \right) \right| , \quad (1.25)$$

and noticing that

$$\left. \begin{aligned} |1-t\lambda u| &\geq 1, 0 \leq t \leq \infty, 0 \leq u \leq 1, \\ 0 \leq \left| u \left(\frac{\gamma}{z} \right) \left(\frac{1-t\lambda uv}{1-t\lambda u} \right) \right| &\leq 1, \quad \left| \gamma/z \right| \leq 1, 0 \leq v \leq 1 \end{aligned} \right\} \quad (1.26)$$

it is easily seen by direct computation that

$$F_n(z, \gamma) \leq M_1^{II} . \quad (1.27)$$

By repeated use of (1.22) and the same method of proof as above, it can be shown that

$$\left. \begin{aligned} F_n(z, \gamma) &\leq M_{q+1}^{II} \\ M_{q+1}^{II} &= \max_{0 \leq w \leq 1} \left| {}_qF_{q+2} \left(\begin{matrix} -n, n+\alpha+\beta+1, \rho_{q-1}, 1 \\ \beta+1, \alpha_{q+1} \end{matrix} \middle| w \right) \right| \end{aligned} \right\} . \quad (1.28)$$

From [2], if $(2\omega + \alpha + \beta + 3/2) > 0$, where

$$2\omega = \left\{ 1/2 + \sum_{i=1}^q \rho_i - \left(q + \beta + \sum_{i=1}^{q+1} \alpha_i \right) \right\}, \quad (1.29)$$

$$M_{q+1}^{II} = \left| \frac{\Gamma(\beta+1)\Gamma(\alpha_{q+1}) N^{4\omega + \alpha + \beta + 1}}{\Gamma(\rho_q - 1)\Gamma(2\omega + \alpha + \beta + 3/2)} \{1 + O(1/N)\} \right| \cdot \left. \begin{array}{l} \\ N^2 = n(n + \alpha + \beta + 1) \end{array} \right\} \quad (1.30)$$

Thus $F_n(z, \gamma)$ is bounded by a term of algebraic order in n .
On the other hand, by (1.18), (1.12) and a result given in [2],

$$\left. \begin{aligned} F_n(\gamma) &= q+3 F_{q+3} \left(\begin{matrix} -n, n+\alpha+\beta+1, \rho_{q-1,1} \\ \alpha_{q+1}, \sigma, 1+\beta \end{matrix} \middle| \gamma/\lambda \right) \\ &= \frac{\Gamma(\alpha_{q+1})\Gamma(\sigma)\Gamma(1+\beta)}{\Gamma(\rho_q-1)2\pi^{1/2}} \left(\frac{N^2 \gamma}{\lambda} \right)^{\bar{\omega}} \exp \left\{ 3 \left(\frac{N^2 \gamma}{\lambda} \right)^{1/3} - \left(\frac{\gamma}{3\lambda} \right) \right\} \\ &+ \frac{\varphi\left(\frac{-\gamma}{\lambda}\right)}{\left(\frac{N^2 \gamma}{\lambda}\right)^{1/3}} + O\left(\frac{\gamma^{7/3}}{N^{4/3}}\right) \left. \begin{array}{l} \\ + \text{terms of algebraic order} \\ \text{in } N \end{array} \right\} \quad (1.31) \end{aligned} \right\}$$

where

$$N^2 = n(n + \alpha + \beta + 1)$$

$$\bar{\omega} = \frac{2\omega + 1/2 - \sigma}{3}$$

$$\bar{\lambda} = -\lambda, \operatorname{Re}(\bar{\lambda}) \geq 0$$

$$\begin{aligned} \varphi(-\gamma/\bar{\lambda}) = & + \frac{1}{15} (\gamma/\bar{\lambda})^2 - (\alpha + \beta + 2/3 + 2\bar{\omega})(\gamma/\bar{\lambda}) \\ & + \frac{(D_1 - D_2)}{3} (2D_1 + D_2 - 1) + E_2 - E_1 - 2/9 \end{aligned} \quad (1.32)$$

$$D_1 = 1 + \sum_{t=1}^q (\rho_t - 1), \quad D_2 = \sum_{i=1}^{q+3} \alpha_i, \quad \alpha_{q+2} = \sigma, \quad \alpha_{q+3} = 1 + \beta$$

$$E_1 = D_1 - 1 + \sum_{n=2}^q \sum_{j=1}^{n-1} (\rho_n - 1)(\rho_j - 1)$$

$$E_2 = \sum_{n=2}^{q+3} \sum_{j=1}^{n-1} (\alpha_n)(\alpha_j)$$

Since $f_n(\gamma)$ is of exponential order in n ,

$$\lim_{n \rightarrow \infty} R_n(z, \gamma) = \lim_{n \rightarrow \infty} \frac{F_n(z, \gamma)}{f_n(z, \gamma)} = 0 \quad (1.33)$$

Similarly in the homogeneous or Type I case, one can show

$$F_n(z, \gamma) \leq \frac{\alpha_{q+1} \lambda^\sigma}{\rho_{q^2}} M_{q+1}^I \quad (1.34)$$

where

$$M_{q+1}^I = \max_{0 \leq w \leq 1} \left| q+3 {}^Fq+2 \left(\begin{matrix} -n, n+\alpha+\beta+1, 1+\rho_{q,1} \\ \beta+1, \alpha_{q+1} \end{matrix} \middle| w \right) \right| \quad \sigma \neq 1$$

$$= \max_{0 \leq w \leq 1} \left| q+3 {}^Fq+2 \left(\begin{matrix} -n, n+\alpha+\beta+1, \rho_{q,2} \\ \beta+1, 1+\alpha_{q+1} \end{matrix} \middle| w \right) \right| \quad \sigma = 1$$
(1.35)

Again, by [2], if $(2n+\alpha+\beta+3/2) > 0$, where

$$2n = \frac{1}{2} + q - \beta + \sum_{i=1}^q \rho_i - \sum_{i=1}^{q+1} \alpha_{q+1}$$

$$= \frac{1}{2} - q - \beta + \sum_{i=1}^q \rho_i - \sum_{i=1}^{q+1} \alpha_{q+1}$$

$$M_{q+1}^I = \left| \frac{\Gamma(\beta+1)\Gamma(\alpha_{q+1}) N^{4n+\alpha+\beta+1}}{\Gamma(1+\rho_q)\Gamma(2n+\alpha+\beta+3/2)} \left\{ 1 + o\left(\frac{1}{N}\right) \right\} \right| \quad \sigma \neq 1$$

$$= \left| \frac{\Gamma(\beta+1)\Gamma(1+\alpha_{q+1}) N^{4n+\alpha+\beta+1}}{\Gamma(\rho_q)\Gamma(2n+\alpha+\beta+3/2)} \left\{ 1 + o\left(\frac{1}{N}\right) \right\} \right| \quad \sigma = 1$$
(1.37)

$$N^2 = n(n+\alpha+\beta+1)$$

Thus, by (1.34) and (1.37), $F_n(z, \gamma)$ is again bounded by a term of algebraic order in n . Combining this with (1.31), we again have

$$\lim_{n \rightarrow \infty} R_n(z, \gamma) = \lim_{n \rightarrow \infty} \frac{F_n(z, \gamma)}{f_n(\gamma)} = 0 \quad (1.38)$$

It calls for remark that the bound for the integral in (1.24) was obtained by ignoring the oscillatory nature of the hypergeometric function in its integrand over the entire region of integration. Thus, the asymptotic bound for the remainder is conservative as the rational approximations converge much more rapidly than indicated by (1.28). Realistic estimates of $F_n(z, \gamma)$ and so of $R_n(z, \gamma)$ seem to be much more difficult. However, though (1.28) is quite rough, it is easy to use and not excessively misleading.

2. Numerics

To give a qualitative idea of the effectiveness of the bound M_{q+1}^{II} for $F_n(z, \gamma)$, we consider an example given in [1] (Table (8.2)). The example is connected with the modified Bessel function $K_\nu(z)$.

$$\left. \begin{aligned} K_\nu(z) &= e^{-z} (\pi/2z)^{1/2} E(z) \\ E(z) &= {}_2F_0(1/2 - \nu, 1/2 + \nu | -1/2z) \end{aligned} \right\} \quad (2.1)$$

Using the non-homogeneous or Type II approach,

$$f_n(\gamma) = {}_3F_3 \left(\begin{matrix} -n, n+\alpha+\beta+1, 1 \\ \beta+1, 1/2-\nu, 1/2+\nu \end{matrix} \middle| -2\gamma \right)$$

$$\Phi_n(z, \gamma) = (1/4 - \nu^2) \sum_{r=0}^n \frac{C_{n,r} (\gamma/z)^r}{(1/2+\nu+r)(1/2-\nu+r)}$$

$$\times {}_3F_3 \left(\begin{matrix} -n+r, n+\alpha+\beta+1+r, 1 \\ \beta+1+r, 3/2+\nu+r, 3/2-\nu+r \end{matrix} \middle| -2\gamma \right)$$

(2.2)

$$E(z, \gamma) = \frac{\Phi_n(z, \gamma)}{f_n(z, \gamma)} + R_n(z, \gamma)$$

$$= E_n(z, \gamma) + R_n(z, \gamma)$$

If $\alpha = \beta = -1/2$, $\nu = 0$, $\gamma = z$, and denoting asymptotic values as computed from (1.30) and (1.31) by \sim , we have to four significant figures,

<u>z</u>	<u>$f_3(z)$</u>	<u>$\sim f_3(z)$</u>	<u>$R_3(z, \gamma)$</u>
0.5	298.3	295.5	5.598×10^{-4}
1.0	1 265.	1 276.	3.145×10^{-4}
2.0	6 515.	7 007.	5.84×10^{-5}
5.0	72 400.	121 000.	0. --
7.0	184 300.	507 800.	$-1. \times 10^{-6}$
10.0	506 600.	3 994 000.	$-8. \times 10^{-7}$

<u>z</u>	<u>$F_3(z, \gamma)$</u>	<u>M_1^{II}</u>	<u>$\sim M_1^{II}$</u>	<u>$\sim M_1^{II} / \sim f_3(z)$</u>
0.5	0.1670	-9.4	9.42	3.188×10^{-2}
1.0	0.3977	↓	↓	7.382×10^{-3}
2.0	0.3805			1.344×10^{-3}
5.0	0.0--			7.785×10^{-5}
7.0	-0.1843			1.855×10^{-5}
10.0	-0.4053			2.359×10^{-6}

(2.3)

<u>z</u>	<u>$f_4(z)$</u>	<u>$\sim f_4(z)$</u>	<u>$R_4(z, \gamma)$</u>
0.5	1 206.	1 198.	1.170×10^{-4}
1.0	7 169.	7 217.	-6.1×10^{-6}
2.0	55 840.	61 170.	-5.4×10^{-6}
5.0	1 208 000.	1 754 000.	$-2. \times 10^{-7}$
7.0	4 052 000.	8 491 000.	$-1. \times 10^{-7}$
10.0	15 140 000.	68 760 000.	$-1. \times 10^{-7}$

<u>z</u>	<u>$F_4(z, \gamma)$</u>	<u>M_1^{II}</u>	<u>$\sim M_1^{II}$</u>	<u>$\sim M_1^{II} / \sim f_4(z)$</u>
0.5	0.1411	12.58	12.57	1.049×10^{-2}
1.0	-0.04373	↓	↓	1.742×10^{-3}
2.0	-0.3015			2.055×10^{-4}
5.0	-0.2417			7.166×10^{-6}
7.0	-0.4052			1.480×10^{-6}
10.0	-1.514			1.828×10^{-7}

(2.4)

The above example shows that our error analysis is quite conservative, as expected. For the present, a pragmatic view should be taken concerning the error. That is, if more precise information is required, in those cases where convergence is assured, one should compute for successive values of n and accept the common digits as correct. To illustrate, in the above example for $K_0(z)$, if

$$E_3(0.5) = 0.85932$$

$$E_4(0.5) = 0.85976$$

and it is quite reasonable to say

$$E(0.5) = 0.860$$

with a possible error of one unit in the third decimal place.

BIBLIOGRAPHY

- [1] Luke, Y. L., "On Economic Representations of Transcendental Functions",
Journal of Mathematics and Physics, 38, No. 4, January 1960, 279-294.
- [2] Fields, J. L., and Luke, Y. L., "Asymptotic Expansions of a Class of
Hypergeometric Polynomials with Respect to the Order", Midwest
Research Institute Technical Report, July 1, 1959.
- [3] Erdélyi, A., Magnus, W., Oberhettinger, F., and Tricomi, F. G.,
Higher Transcendental Functions, Vols. I and II, McGraw-Hill Book
Company (1953).
- [4] Luke, Y. L., "Remarks on the \mathcal{Z} -Method for the Solution of Linear
Differential Equations with Rational Coefficients", SIAM, 3 (1955),
179-191.

DISTRIBUTION LIST

Contract No. Nonr-2638(00)(X)

No. of Copies

Recipient

14

CHBUSHIPS

10 Tech. Library (Code 312)

1 Tech. Asst. to Chief (Code 106)

1 Electronic Computer Div. (Code 280)

1 Asst. Chief for Field Activities
(Code 700)

1 Asst. Chief for Nuclear Propulsion
(Code 1500)

1

CHBUAER

1

CHBUORD

1

CHBUSANDA

1

CHBUCENSUS

1

CHONR

1

NAVSHIPYD BSN

1

NAVSHIPYD CHASN

1

NAVSHIPYD LBEACH

1

NAVSHIPYD MARE

1

NAVSHIPYD NYK

1

NAVSHIPYD NORVA

1

NAVSHIPYD SFRAN

1

NAVSHIPYD PHILA

1

NAVSHIPYD PTSMH

1

NAVSHIPYD PUG

1

NAVSHIPYD PEARL

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	CO & DIR, USNBTL Philadelphia, Pennsylvania
1	CO & DIR, USNEL San Diego, California
1	CO & DIR, USNRDL San Francisco, California
1	CO & DIR, USN Trg. Device Ctr. Computer Branch Port Washington, New York
1	CO, USNCML St. Paul, Minnesota
1	CDR, USNPG Dahlgren, Virginia
3	CDR, USNOTS China Lake, California 1 Pasadena Annex 1 Tech. Library 1 Michelson Lab (Code 5038)
1	CDR, USNOL White Oak, Silver Spring, Maryland
1	DIR, USNEES Annapolis, Maryland
1	DIR, USNRL Washington 25, D. C.
1	SUPT, USN Postgrad School Monterey, California Attn: Library, Technical Reports Section
1	SUPT, US Naval Academy Department of Math Annapolis, Maryland

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	CG, Aberdeen Proving Ground Aberdeen, Maryland
1	DIR, National Bureau of Standards Washington 25, D. C.
1	Chief, AFSWP Washington 25, D. C.
1	DIR, Langley Aero Lab Langley Field, Virginia
1	DIR, Lewis Flight Propulsion Lab NACA Cleveland 11, Ohio
1	CDR, Wright Air Development Center Attn: WCRRN-4 Wright-Patterson Air Force Base, Ohio
1	CG, Frankford Arsenal Head, Math Section Pitman-Dunn Laboratories Philadelphia 37, Pennsylvania
1	CG, White Sands Proving Ground Flight Determination Lab Las Cruces, New Mexico
1	USAEC, Technical Information Service Oak Ridge, Tennessee
2	U. S. Atomic Energy Commission Washington 25, D. C. Attn: Tech. Library
1	CO, Diamond Ordnance Fuze Laboratory Washington 25, D. C. Attn: Library
1	George Washington University Logistics Research Washington, D. C.

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Johns Hopkins University Applied Physics Laboratory Silver Spring, Maryland
1	University of California Librarian, Numerical Analysis Los Angeles 24, California
1	Carnegie Institute of Technology Pittsburgh, Pennsylvania
1	Hudson Laboratory Columbia University Dobbs Ferry, New York
2	Harvard University Cambridge, Massachusetts 1 Department of Math Attn: Prof. J. L. Walsh 1 Computation Laboratory
1	Institute for Advanced Study Princeton, New Jersey
2	Massachusetts Institute of Technology Cambridge, Massachusetts Attn: Computation Center
2	New York University New York, New York 1 Inst. of Math Sciences 1 AEC Computing Facility
1	DIR, Research Center Ohio State University Columbus, Ohio
1	Pennsylvania State University Department of Math University Park, Pennsylvania

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Princeton University Princeton, New Jersey Attn: Library
1	University of California Berkeley, California
2	University of Illinois Urbana, Illinois 1 Department of Math 1 Electronic Digital Computer Project
2	University of Maryland College Park, Maryland 1 Department of Math 1 Inst. for Fluid Dynamics and Applied Math
1	Willow Run Research Center University of Michigan Ypsilanti, Michigan
1	University of Washington Department of Math Seattle, Washington
1	Yale University New Haven, Connecticut
1	State College of Washington Department of Math Pullman, Washington
1	Stanford University Palo Alto, California Attn: Dr. G. E. Forsythe
1	Johns Hopkins University Charles and 34th Streets Baltimore 18, Maryland
1	Rutgers University New Brunswick, New Jersey Attn: Prof. E. P. Starke

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Brown University Division of Engineering Providence, Rhode Island Attn: Dr. R. D. Kodis
1	Princeton University Princeton, New Jersey Attn: Prof. H. J. Maehly Chief of Computer Project
1	University of Rochester Department of Math Rochester, New York
1	The Scripps Institute of Oceanography University of California LaJolla, California Attn: Dr. Walter Munk
1	Dr. Werner C. Rheinboldt, Director Computing Center, 112 Hinds Hall Syracuse University Syracuse 10, New York
1	DIR, Combustion Engineering, Inc. Windsor, Connecticut
1	DIR, Westinghouse Electric Corporation Bettis Atomic Power Division P.O. Box 1468 Pittsburgh 30, Pennsylvania
2	Argonne National Laboratory P.O. Box 299 Lemont, Illinois
1	Armour Research Foundation 35 W. 33rd Street Chicago 16, Illinois
1	Battelle Memorial Institute 505 King Avenue Columbus, Ohio

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Brookhaven National Laboratory Upton, Long Island, New York
1	Cornell Aero Laboratory, Inc. Buffalo, New York
1	Curtiss-Wright Corporation Research Division Clifton, New Jersey
1	Douglas Aircraft Company, Inc. Santa Monica Division 3000 Ocean Park Boulevard Santa Monica, California
1	Engineering Research Associates St. Paul 4, Minnesota
1	IBM Corporation 590 Madison Avenue New York 22, New York
1	Knolls Atomic Power Laboratory General Electric Company Math Analysis Unit Schenectady, New York
1	Lincoln Lab B-125 Lexington, Massachusetts
1	Lockheed Aircraft Corporation Missile Systems Division Sunnyvale, California
1	Lockheed Aircraft Corporation Van Nuys, California
1	Los Alamos Scientific Lab Los Alamos, New Mexico

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Remington Rand Division of Sperry Rand Electronic Computer Department 315 Fourth Avenue New York 10, New York
1	Ramo-Wooldridge Corporation 8820 Bellanca Avenue Los Angeles 45, California
1	Rand Corporation Santa Monica, California
1	Sandia Corporation Albuquerque, New Mexico Attn: Library
1	United Aircraft Corporation 400 Main Street East Hartford 8, Connecticut
1	Vitro Corporation of America 261 Madison Avenue New York 16, New York
1	Oregon State College Corvallis, Oregon Attn: Professor W. Milne
1	Professor A. C. Aitken University of Edinburgh Edinburgh, Scotland
1	Professor D. E. Rutherford University of St. Andrews St. Andrews, Scotland
1	Professor J. L. Synge Dublin Institute for Advanced Studies Dublin, Eire

DISTRIBUTION LIST (Continued)

<u>No. of Copies</u>	<u>Recipient</u>
1	Professor T. S. Broderick Trinity College Dublin, Eire
1	Professor R. E. Langer Mathematics Research Center University of Wisconsin Madison 6, Wisconsin
1	Professor Wallace Givens Department of Mathematics Wayne State University Detroit 2, Michigan
1	Dr. A. S. Householder Oak Ridge National Laboratories Oak Ridge, Tennessee
1	Professor John Todd Department of Math California Institute of Technology Pasadena, California
1	Dr. Franz Alt Mathematical Computation Laboratory National Bureau of Standards Washington 25, D. C.
1	Dr. E. W. Cannon Applied Mathematics Division National Bureau of Standards Washington 25, D. C.
1	Miss Irene Stegun Mathematical Computation Laboratory National Bureau of Standards Washington 25, D. C.
1	Mr. Cecil Hastings, Jr. 136 Kunalu Street Kailua, Hawaii

DISTRIBUTION LIST (Concluded)

<u>No. of Copies</u>	<u>Recipient</u>
1	Professor E. J. McShane University of Virginia Charlottesville, Virginia
1	Professor J. B. Rosser Department of Math Cornell University Ithaca, New York
1	Professor D. H. Lehmer Department of Math University of California Berkeley 4, California
1	Professor Arthur Erdélyi Department of Math California Institute of Technology Pasadena, California
5	Navy Department David W. Taylor Model Basin Washington, D. C. Attn: Dr. John W. Wrench, Jr.
1	Professor Bertram Bussell Department of Engineering University of California Los Angeles 24, California
75	Midwest Research Institute Attn: Yudell L. Luke
1	Midwest Research Institute Attn: Mathematics and Physics Division
1	Midwest Research Institute Attn: Library